

OPEN ACCESS

Microbiology

Research Article

Improvement of paromomycin production by *Streptomyces rimosus* subsp *paromomycinus* NRRL 2455 using gamma irradiation mutagenesis

Asmaa A. Ibrahim, Khaled M. Aboshanab^{*}, Mahmoud A. Yassien, Nadia A. Hassouna

Department of Microbiology and Immunology, Faculty of Pharmacy, Ain Shams University, Cairo, Egypt

ABSTRACT

Streptomyces (S.) rimosus NRRL 2455 produces paromomycin, a 2-deoxystreptamine aminocyclitol aminoglycoside antibiotic (2DOS-ACAGA) with broad-spectrum activity against most of the Gram-positive and Gram-negative bacteria as well as protozoa. The mutation has become one of the beneficial methods used in enhancing the microbial production. Improvement of the paromomycin production by *S. rimosus* NRRL 2455 was achieved via irradiation mutagenesis using gamma (γ) radiation. The culture of *S. rimosus* was irradiated with different doses of gamma radiation (3, 4 and 5 KiloGray (KGy) to find out the best dose for the mutation that gave 99.99% killing. The optimum dose was found to be 4 KGy. Six morphologically changed colony types appeared on tryptic soy agar plates. These colonies were bio-assayed for their antimicrobial activity against standard *Staphylococcus aureus* ATCC 25923 using agar well diffusion technique. A mutant coded 5M showed about 1.44, and 2 fold increase in its activity as compared with the wild-type when cultivated in basal culture or optimized media (soybean meal 30 g/L, NH₄CL 4 g/L, CaCO₃ 5 g/L and glycerol 40 ml/L), respectively. Moreover, high genetic stability was observed upon subsequent culturing of 5M-mutant. Therefore, *S. rimosus* mutant-5M can be used as a potential industrial strain for paromomycin production.

Keywords: Paromomycin, *Streptomyces rimosus*, gamma mutagenesis, 2DOS-ACAGA.

***Correspondence** | Khaled M. Aboshanab; Professor of Microbiology and Immunology, Faculty of Pharmacy, Ain Shams University, African union organization street Abassia, 11566, Cairo, Egypt. Email: aboshanab2012@pharma.asu.edu.eg

Citation | Asmaa AI, Khaled MA, Mahmoud AY, and Nadia AH. 2017. Improvement of paromomycin production by *Streptomyces rimosus* subsp *paromomycinus* NRRL 2455 using gamma irradiation mutagenesis. Arch Pharm Sci ASU 1(2): 26-30

DOI: [10.21608/aps.2017.11017](https://doi.org/10.21608/aps.2017.11017)

Online ISSN: 2356-8380

Print ISSN: 2356-8399

Copyright: © 2017 Ibrahim et al. This is an open-access article licensed under a Creative Commons Attribution 4.0 International License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

Published by: Ain Shams University, Faculty of Pharmacy.

1. INTRODUCTION

Streptomyces is a member of the family Actinomycetaceae. It is characterized by having mycelia and spores at the aerial hypha. They are attributed to the production of secondary metabolites such as antibiotics [1]. *S. rimosus* NRRL-2455 produces paromomycin, a 2 deoxystreptamine (2DOS)-containing aminoglycoside antibiotic with broad-spectrum activity against Gram-negative bacteria and most of the Gram-positive bacteria especially *Staphylococcus* strains particularly those resistant to oxytetracycline, erythromycin or carbomycin

[2,3]. Members of *Streptomyces* display genetic instability, intra-strain morphological differences and there is also a correlation between colony morphology and antibiotic activity [4]. Random mutagenesis is considered as an effective way to improve the productivity of industrial microbial cultures [5]. It was found that some mutants with altered colony morphology and resulting from basic genetic studies exhibited enhanced activities [6]. Also, the change in the medium composition affects greatly the biosynthesis of the produced antibiotic from streptomycetes [7]. The current study aimed at enhancing the

paromomycin production from *S. rimosus* subsp. *paromomycinus* NRRL 2455 through gamma irradiation mutagenesis and showing the effect of the wild strain optimized media (unpublished data) on the hyper producer mutant.

2. MATERIALS AND METHODS

2.1. Microorganisms

S. rimosus subsp. *paromomycinus* NRRL 2455 was stored in tryptic soy broth (TSB) containing 50% glycerol at -20 °C [8,9] and cultured in TSB at 28 °C, 200 rpm and a 3-day incubation period for stock preparation to be used later for the preparation of a seed culture. *Staphylococcus aureus* ATCC 25923 was stored in Luria–Bertani broth (LB broth) [10] containing 20% glycerol at -20°C and cultured on nutrient agar slants at 37 °C for 24 h to be used in the bioassay of paromomycin production using agar well diffusion technique.

2.2. Culture Media

The basal medium for *S. rimosus* subsp. *paromomycinus* and the mutants were TSB [9]. *S. rimosus* optimized media was composed of soybean meal 30 g/L, NH₄CL 4g/L, CaCO₃ 5g/L and glycerol 40 ml/L.

2.3. Preparation of bacterial culture for paromomycin production

The seed culture was prepared by inoculating 25 ml basal medium in a 250 mL Erlenmeyer flask with 1.375 ml of the prepared stock and incubated for 72 h at 28 °C, 200 rpm. The main culture of *S. rimosus* was done in 25 mL TSB in 250 mL Erlenmeyer flasks at 28 °C, 200 rpm, and was then inoculated from the prepared seed culture.

2.4. Mutation Using Gamma Irradiation

After five days of incubation five mL aliquots from the above mentioned main culture (1.5×10^7 CFU/mL) were irradiated with different doses of gamma radiation (3, 4 and 5 KiloGray (KGy)) to find out the best dose for a mutation that gave 99.99% killing. The source of gamma radiation was ⁶⁰Co from Indian Gamma cell that was

providing a dose rate of 2.2 KGy/h at the time of the experiment. The irradiation process was carried out at The National Center for Radiation Research and Technology, Atomic Energy Authority, Nasr City, Cairo, Egypt. After mutagenesis, the irradiated cell suspension was suitably diluted, plated on tryptic soy agar plates and incubated at 28 °C to get single colonies.


2.5. Determination of the antibacterial activity of the obtained mutants

To examine the antibacterial activity of the paromomycin produced by *S. rimosus* subsp. *paromomycinus* and its mutants, every single colony of the mutants and wild-type were conveyed to 25 mL of fresh TSB in a 250 mL Erlenmeyer flask and incubated for 3 days at 28 °C and 200 rpm. The obtained broths were used for the inoculation of the seed culture which was then used to inoculate the main culture as well as *S. rimosus* optimized media as the same procedure mentioned above. After each incubation process, the culture broths were centrifuged and sterilized by filtration using 0.22 µm pore size cellulose membrane filters (CHEM LAB, Barcelona, Spain). The culture filtrates were bio-assayed against *Staphylococcus aureus* ATCC 25923 by using agar well diffusion technique [11]. The suspension (0.5 McFarland) of *Staphylococcus aureus* ATCC-25923 was uniformly and aseptically spread on Mueller Hinton agar surface (MHA, Difco, USA) and 150 µl of the culture filtrate was used to fill the wells. Plates were kept at 4-8 °C for at least 30 min to allow the diffusion of culture filtrate. The diameters of inhibition zones were recorded after 24 h of incubation at 37 °C. All the experiments were done in triplicate and the mean of the three readings was documented.

3. RESULTS

The optimum dose of gamma radiation for the mutation that causes 99.99% killing was found to be 4 KGy. Gamma irradiation of wild-type strain caused the formation of six morphologically changed colony types, coded (1M, 2M, 3M, 4M, 5M, and 6M), on tryptic soy agar plates. These colonies were bio-assayed for their antimicrobial activity as shown in **Fig. 1**. All the mutants


obtained showed higher antimicrobial activities than the wild type except for the two mutants 3M



and 4M.


Fig. 1 Inhibition zone diameters of the six morphologically changed Mutants and Wild-type *S. rimosus* in the agar well diffusion assay.

The morphology of the mutant coded 3M colony showed the absence of the aerial mycelia as shown in **Fig. 2** and this morphological change was associated with loss of the antimicrobial



activity.

Fig. 2 The morphology of Mutants 5M and 3M compared to



wild-type on tryptic soy agar plates.

Fig. 3 Inhibition zone diameters of the wild-type and Mutant 5M after culturing in basal media and optimized media.

The best mutant was mutant 5M which showed about 1.44 and 2 fold increase in its activity as compared to the wild-type when

cultivated in basal culture or optimized media, respectively as shown in **Fig. 3** Furthermore, successive culturing of 5M-mutant exhibited high genetic stability.

4. DISCUSSION

The mutation has been used for strain development to enhance the beneficial microbial secondary metabolites yield produced⁵ and the screening for overproducing microbial mutants. These mutants' strains being overproducers offer the advantage of minimizing the production costs [6]. The most widely used mutagenic agents include methyl methanesulfonate (MMS), hydroxylamine (HA), ethyl methanesulfonate (EMS), N-methyl-N-nitro- N-Nitroso guanidine (MNNG) and ultraviolet (UV) irradiation [12]. The improvement of actinomycetes mutants has long been realized. The exposure of *Streptomyces* spores to ultraviolet and gamma rays develops hereditary changes that consequently cause colony morphology and pigmentation. These changes are greatly related to instabilities. These instabilities generate novel variants [13]. It is known that *Streptomyces* DNA contains high G+C content [14]. Furthermore, the best way to yield improvement is AT to CG transversions. So far, no AT to CG transversions has been reported by any chemical mutagen [12]. On the other hand, AT to CG transversions were found in cells treated with gamma rays [15]. Gamma rays induced mutation occurs through single or double-strand breakage of DNA resulting from deletion or structural change [16]. The application of gamma irradiation and the chosen dose depends on the types of biological materials where high doses are used for sterilization, medium doses for decontamination while low doses are used for mutagenesis [17]. In our study, *S.rimosus* subsp. *paromomycinus* was subjected to gamma rays producing six morphologically changed colonies. These colonies were bio-assayed against standard *Staphylococcus aureus* ATCC 25923. A mutant coded 5M showed about 1.44, and 2 fold increase in its activity as compared to the wild-type when cultivated in basal culture or optimized media respectively. Similar results were reported in previous studies. *Aspergillus niger*, a potent producer of many

important industrial enzymes, was genotypically improved by exposure to gamma rays [18]. *Aspergillus niger* producing α - and β -galactosidases enzymes revealed a two-fold increase in these two enzymes produced by its mutant after exposure to gamma radiation [19]. Also enhanced production of glucose oxidase was reported by mutants of *Aspergillus niger* after a dose of 80 krad [18], while at a dose of 2 KGy, Mutant of *Aspergillus niger* improved its carboxymethylcellulose and filter paper cellulase production [20]. Moreover, Shahbazi et al. (2014) stated that a mutant strain of *Trichoderma reesei* irradiated at 250 Gy of gamma ray (PTCC 5142) showed the maximum production of cellulase enzyme compared with UV-irradiated mutant and wild-type strains [21]. The tremendous increase was also reported with oxytetracycline production from a *Streptomyces rimosus* CN08 mutant which was called *Streptomyces rimosus* γ -45 [22]. Besides, a 12 fold increase was reported in the production of epsilon-pyrromycinone glycosides in *Streptomyces galilaeus* with respect to the wild strain after both natural selection and treatment with mutagens as gamma-irradiation [23]. Additionally in our findings, Successive culturing of the mutant 5M revealed great genetic stability. Similar results were reported for *Streptomyces fradiae* NRRL-2702 mutant gamma-1, however, UV irradiation associated changes were unstable with loss of tylosin activity [24]. These results make gamma radiation an effective way for mutation compared to other mutagens.

5. Conclusion

Genetic manipulation was carried out using gamma radiation and results revealed that gamma irradiation mutagenesis could generate production associated morphological changes in *S. rimosus* NRRL-2455. It was also noted that subsequent replating of the mutants showed same morphological changes and also constant paromomycin production upon successive culturing. A mutant coded 5M, the best mutant, showed about 1.44 increases in its activity as compared to the wild-type when cultivated in basal culture. It was interesting that culturing of mutant 5M in the *S. rimosus* subsp

paromomycinus optimized media revealed a 2 fold increase in the production which is promising for scaling up the paromomycin production from this mutant strain. Therefore, Gamma Radiation was found to be very effective for improvement of paromomycin production and the obtained *S. rimosus* mutant-5M may be used as a potential industrial strain for paromomycin production in future studies.

Acknowledgment

We hereby acknowledge the Department of Microbiology and Immunology, Faculty of Pharmacy, Ain Shams University for providing us with all the facilities and support required to perform the practical work.

REFERENCES

1. Sharma A, Gautam S, Saxena S. *Streptomyces*. In: Batt CA, Tortorello ML, eds. Encyclopedia of Food Microbiology (Second Edition). Oxford: Academic Press, 2014:560–566.
2. Aboshanab KMA. Isolation, recombinant expression and characterization of the dprA gene product in *Streptomyces rimosus* NRRL 2455. African Journal of Microbiology Research 2010; 4(10):915–922.
3. Frohardt RP, Haskell TH, John E, Penner KM. Antibiotic and methods for obtaining the same 1959.
4. Schrempf H. Plasmid loss and changes within the chromosomal DNA of *Streptomyces reticuli*. Journal of Bacteriology 1982; 151(2):701–707.
5. Parekh S, Vinci VA, Strobel RJ. Improvement of microbial strains and fermentation processes. Appl Microbiol Biotechnol. 2000; 54(3):287–301.
6. Adrio JL, Demain AL. Genetic improvement of processes yielding microbial products. FEMS Microbiol Rev 2006; 30(2):187–214.
7. Abbanat D, Maiese W, Greenstein M. Biosynthesis of the pyrroindomycins by *Streptomyces rugosporus* LL-42D005; characterization of nutrient requirements. The J Antibiot 1999; 52(2):117–126.
8. Hopwood DA, Wright HM. Bacterial protoplast fusion: recombination in fused protoplasts of *Streptomyces coelicolor*. Molecular & general genetics: MGG 1978; 162(3):307–317.

9. Kieser T, Bibb M, Buttener M, Chater K, Hopwood D. Practical *Streptomyces* Genetics, 2000.
10. Miller JH. Experiments in molecular genetics. Cold Spring Harbor Laboratory, 1972.
11. Balouiri M, Sadiki M, Ibsouda SK. Methods for in vitro evaluating antimicrobial activity: A review. J Pharm Anal 2016; 6(2):71–79.
12. Baltz RH. Genetic methods and strategies for secondary metabolite yield improvement in actinomycetes. Antonie van Leeuwenhoek 2001; 79(3–4):251–259.
13. Guravaiah M. Production of alkaline protease by *Streptomyces indicus*. India: DARSHAN PUBLISHERS, 2016.
14. Surajit D, Lyla PS, Khan seyed. Distribution and generic composition of culturable marine actinomycetes from the sediments of Indian continental slope of Bay of Bengal. Chinese Journal of Oceanology and Limnology 2008; 26(2):166-177
15. Xie, An X, Li-Jun W, Jian-Min Y, Jian-Bo Y, Zeng-Liang Y. Comparison of base substitutions in response to nitrogen ion implantation and ⁶⁰Co-gamma-ray irradiation in *Escherichia coli*. Genet Mol Biol 2004; 27(2):284–290.
16. Huma T, Rashid MH, Javed MR, Ashraf A. Gamma-ray mediated mutagenesis of *Phialocephalohumicola*: Effect on kinetics and thermodynamics of α-amylase production. African Journal of Microbiology Research 2012; 6(22):4639–4646.
17. Hoe PCK, Rahim KA, Saud HM. A Review on Microbial Mutagenesis through Gamma Irradiation for Agricultural Applications. Jurnal Sains NuklearMalaysia 2016; 28(2):20–29.
18. Zia MA, Rasul S, Iftikhar T. Effect of gamma irradiation on *Aspergillusniger* for enhanced production of glucose oxidase 2012; 44(5).
19. Awan MS, Tabbasam N, Ayub N, Babar ME, Mehboob-ur-Rahman null, Rana SM, Rajoka MI. Gamma radiation-induced mutagenesis in *Aspergillusniger* to enhance its microbial fermentation activity for industrial enzyme production. Molecular Biology Reports. 2011; 38(2):1367–1374.
20. Mostafa A. Effect of gamma irradiation on *Aspergillusniger*DNA and production of cellulases enzymes The Journal of American Science 2014; 10(5):152–160.
21. Shahbazi S, Ispareh K, Karimi M, Askari H, Ebrahimi M. Gamma and UV radiation-induced mutagenesis in *Trichoderma reesei* to enhance cellulases enzyme activity. International Journal of Farming and Allied Sciences 2014; 3(5):543–554.
22. Lazim H, Slama N, Mankai H, Barkallah I, Limam F. Enhancement of oxytetracycline production after gamma irradiation-induced mutagenesis of *Streptomyces rimosus* CN08 strain. J Microbiol Biotechnol 2010; 26(7):1317–1322.
23. Královocová E, Blumauerová M, Vaněk Z. Strain improvement in *Streptomyces galilaeus*, a producer of anthracycline antibiotics galirubins. Folia Microbiologica 1977; 22(5):321–328.
24. Khaliq S, Akhtar K, AfzalGhauri M, Iqbal R, Mukhtar Khalid A, Muddassar M. Change in colony morphology and kinetics of tylosin production after UV and gamma irradiation mutagenesis of *Streptomyces fradiae* NRRL-2702. Microbiological Research 2009; 164(4):469–477.