

The protective role of some natural antioxidants against some nanoparticles-induced subchronic nephrotoxicity in Wistar rats

Riyadh MUSAED Naji^{a,b}, Abdallah H. Fathy^{c*}, Mohamed A. Bashandy^b

^aDepartment of Zoology, Faculty of Science, Al-Azhar University, Cairo 11651, Egypt

^bDepartment of Zoology, Faculty of Science and Education, Aden University, Yemen

^cDepartment of Animal House Facility, Faculty of Pharmacy, Ain Shams University, Cairo 11566, Egypt

ABSTRACT

The aim is to study the possible protection of fig fruit extract with olive oil and date-palm fruit extract (FOD) as natural antioxidants in decreasing the subchronic toxicity hazards of silicon oxide nanoparticles (SiO₂NPs), aluminum oxide nanoparticles (Al₂O₃NPs), or zinc oxide nanoparticles (ZnONPs) in male rats treated for 75 days. We used 80 male Wistar rats distributed into 8 groups (n = 10) according to the experiment. We used various antioxidant treatments at their recommended antioxidant doses. All nanoparticles were given orally and daily for 75 days at doses of 100 mg/kg. The oral administration of different nanoparticles (NPs) alone led to dramatic histopathological features, a significant increase in the levels of the TBARS, tumor suppressor p53, and inflammatory markers (TNF- α and IL-6) in the renal tissue. In addition, serum kidney function parameters elevated significantly in NPs treated groups compared with the control group. On the other hand, the renal TAC, GSH, SOD, and TBARS lowered significantly in the renal tissue of rats administered with different NPs compared to the control group. The FOD-NPs-treated groups recorded significantly reduced nephrotoxicity effects as compared to the groups treated with NPs alone. In Conclusion, administration of FOD provides considerable protective effects against NPs-induced subchronic nephrotoxicity in male Wistar rats.

Keywords: SiO₂NPs; Al₂O₃NPs; ZnONPs; olive oil; extracts; oxidative stress; P53; kidney.

*Correspondence | Abdallah H. Fathy; Animal House Facility, Faculty of Pharmacy, Ain Shams University, Cairo 11566, Egypt.

Email: abdallahhfathy@pharma.asu.edu.eg; abdallah.ahf@gmail.com

Citation | Naji RM, Fathy AH, Bashandy MA, 2022. The protective role of some natural antioxidants against some nanoparticles-induced subchronic nephrotoxicity in Wistar rats. Arch Pharm Sci ASU 6(1): 1-16

DOI: [10.21608/aps.2022.124160.1082](https://doi.org/10.21608/aps.2022.124160.1082)

Print ISSN: 2356-8380. **Online ISSN:** 2356-8399.

Received 26 February 2022. **Accepted** 25 March 2022.

Copyright: ©2022 Naji *et al.* This is an open-access article licensed under a Creative Commons Attribution 4.0 International License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

Published by: Ain Shams University, Faculty of Pharmacy

1. INTRODUCTION

Nanotechnology creates products that exhibit novel properties (particle size range from 1 to 100 nm) [1]. The spread of nanotechnology in industry causes the possible accumulation of nanomaterials in the human body exposed to these NPs [2]. Nanoparticles affect various physiological systems and may enter the human body via various routes and may even cause

pathological disorders [3].

The applications of SiO₂NPs in industry and biomedicine products are promising because they possess stabilities and easy modifications [4]. Toxicology studies have suggested that SiO₂ NPs can induce adverse effects in the liver, kidney, and brain [5-7]. Al₂O₃NPs have been used in industry [8]. The acute exposure to Al₂O₃NPs causes toxicity in experimental animals [9]. ZnONPs are used in food additives and other

pharmaceuticals [10, 11]. Subsequently, the increased use of ZnONPs causes more concerns about their toxicity and inflammatory effects [12].

The Mediterranean diet contains natural products that produce a promising effect on the health state [13]. *Ficus carica* (fig) fruits (belonging to the Moraceae family), typically Mediterranean species have antioxidant properties. The fig fruits contain high levels of polyphenols, flavonoids, and anthocyanins [14]. Moreover, fig-trees possess multiple therapeutic and anti-inflammatory activities against different toxins [15]. Olive oil is an important constituent of the Mediterranean diet. It contains many antioxidants and active compounds responsible for its biological activities as oleic acid, phenolics (including hydroxytyrosol, tyrosol, and oleuropein), and squalene [16]. *Phoenix dactylifera* (belonging to the family Arecaceae) is one of the species of date palm and is considered one of the oldest cultivated fruit trees in the Middle East. The flesh of the dates contains a higher percentage of carbohydrates, vitamin C, vitamin A, vitamin E, vitamin B2, and dietary fibers. In addition, it is rich in iron, calcium, magnesium, phosphorus, potassium, sodium, zinc, and selenium and protects against oxidative stress [17-19].

This study aims to explore the possible protective effects of FOD against different nanoparticles-induced subchronic nephrotoxicity in Wistar rats including renal oxidative stress, some biochemical, inflammatory, and histopathological alterations in addition to P53 content in the kidney tissue with a view to its possible applications in the clinical field.

2. MATERIALS AND METHODS

2.1. Chemicals

We used chemicals and reagents of high analytical grade from standard suppliers.

2.2. Nanoparticles

The silicon oxide nanoparticles (SiO₂NPs), aluminum oxide nanoparticles (Al₂O₃NPs), or zinc oxide nanoparticles (ZnONPs) were prepared and characterized in a private laboratory (Nanogate Laboratory, Cairo, Egypt).

2.3. Preparation of nanoparticles (NPs) treatments

The nanoparticles (SiO₂NPs, Al₂O₃NPs, and ZnONPs) were suspended in water. This suspension was vibrated by vortex for 5 min before injection to aid in preparing a homogeneous suspension. All nanoparticles were given orally by oral gavages for 75 consecutive days. All nanoparticles (SiO₂NPs [20], Al₂O₃NPs [21], and ZnONPs [22]) were given at doses of 100 mg/kg according to pilot studies in our lab (data not shown) and the given doses were found following previously published studies.

2.4. Plant materials and authorities

We purchased the extra-virgin olive oil from the Grup Pons Company (Spain), the fig fruit from Kafods Ltd. (Turkey), and the date-palm fruit from the Al-MADINA AL-MUBARAK market (Saudi Arabia).

2.5. Preparation of crude extracts

Ficus carica fruit extract was prepared and lyophilized according to a previous method [23]. The hydroalcoholic extract of the date fruit was made and lyophilized according to a previous method [24].

2.6. Preparation of the antioxidant treatments

The extra-virgin olive oil was supplemented to rats by oral gavage. Every rat received oil in a concentration of 7 g/kg body mass [25].

The fig and date-palm fruit extracts were prepared for supplementation in rats at a concentration of 1 g/mL just before the experimental use. The selected fig and date-palm

fruit doses were calculated based on the human recommended antioxidant doses [26] after conversion to rat doses.

2.7. The experimental animals

The experimental animals used in the present study were the mature male Wistar albino rats obtained from the Egyptian Holding Company for Biological Products and Vaccines (VACSERA, Giza, Egypt) and were housed at the animal facility, Faculty of Pharmacy, Ain Shams University, Cairo, Egypt and provided with diet pellets and tap water. Five rats were placed into each cage and provided with standard diet pellets and drinking tap water *ad libitum* during the experimental period. Animals in all experiments remained 2-week acclimatization

Table 1. The study design

Groups	Treatment	Period
1. Control	diet pellets and tap water	2 W + 75 days
2. FOD	extra-virgin olive oil (7 g/kg) and freshly prepared fig extract (1 g/kg) and date-palm fruit extract (1 g/kg) daily and orally.	2 W + 75 days
3. SiO ₂ NPs	SiO ₂ NPs in doses of 100 mg/kg, daily and orally.	75 days
4. FOD-SiO ₂ NPs	FOD and SiO ₂ NPs on the same schedule mentioned above in groups II and III.	2 W + 75 days
5. Al ₂ O ₃ NPs	Al ₂ O ₃ NPs in doses of 100 mg/kg, daily and orally.	75 days
6. FOD-Al ₂ O ₃ NPs	FOD and Al ₂ O ₃ NPs on the same schedule mentioned above in groups II and V.	2 W + 75 days
7. ZnONPs	ZnO in doses of 100 mg/kg, daily and orally.	75 days
8. FOD-ZnONPs	FOD and ZnONPs on the same schedule mentioned above in groups II and VII.	2 W + 75 days

Note: FOD, fig with olive oil and date-palm fruit extracts; SiO₂NPs, silicon oxide nanoparticles; Al₂O₃NPs, aluminum oxide nanoparticles; ZnONPs, zinc oxide nanoparticles; W; week

period before treatments. In addition, the animal masses were recorded regularly one time per week and the animal behaviors were monitored daily.

2.8. Animal welfare

The *in vivo* studies were conducted under the National Research Centre guidelines for the use and care of laboratory animals [27] and were approved by an independent ethics committee of the Faculty of Pharmacy, Ain Shams University.

2.9. Experimental design

A patch of 80 male Wistar albino rats average weight of 150-170 g were allowed to acclimatize for 2 weeks, and then divided into 8 groups as represented in **Table 1**

2.10. Collection of samples

Samples were collected from each animal under anesthesia from the retro-orbital venous plexus puncture at the end of the experiment. Blood samples were collected in non-heparinized tubes and centrifuged at 4000 rpm for 10 min. The sera were frozen at -80 °C for the following measurements. After blood sampling, the animals were dissected and the kidneys were quickly removed and washed with saline, and divided into two portions one was used for biochemical analysis and the other was used for histopathology. The portion used for biochemical analysis was homogenized in ice-cold Tris-HCl lysis buffer, pH 7.4 using rotor-stator Homogenizer, fitted with a Teflon pestle (Omni International, Kennesaw, GA, USA). The homogenates were centrifuged under cooling at 3000 rpm for 20 min. The supernatants were aliquoted and stored at -80 °C until used.

2.11. Biochemical study

2.11.1. Renal oxidative stress parameters

The measurements of oxidative stress markers of the renal tissue (renal total antioxidant capacity (TAC), renal reduced glutathione (GSH), and renal superoxide dismutase (SOD), thiobarbituric acid reactive substances (TBARS)) were estimated in the kidney tissue homogenate using kits from bio-diagnostic Co. for research kits, Egypt.

2.11.2. Inflammatory markers

The tumor necrosis factor-alpha (TNF- α) was estimated in the kidney tissue homogenate of rats by ELISA technique using a kit purchased from MyBioSource, Inc. San Diego, CA 92195-3308: USA according to the manufacturer's instructions provided with the TNF- α assay kit (Catalog No: MBS2507393). The interleukin-6 (IL-6) was estimated in the kidney tissue homogenate of rats by ELISA technique using a kit purchased from Elabscience Biotechnology, Inc., Texas, USA

according to the manufacturer's instructions provided with the IL-6 assay kit (Catalog No: E-EL-R0015).

2.11.3. Apoptotic Biomarkers Estimation

The P53/Tumor Protein (p53/TP53) was estimated in the kidney tissue homogenate of rats by ELISA technique using a kit purchased from Cusabio Technology LLC, Houston, USA using methods outlined in the diagnostic kit.

2.11.4. Kidney function tests

The measurements of serum kidney function parameters (creatinine, urea, blood urea nitrogen (BUN), and uric acid) were estimated in the blood serum using kits from bio-diagnostic Co. for research kits, Egypt.

2.11.5. The histopathological study

The kidney tissue samples were fixed, processed, sectioned, and stained according to certain methods [28].

2.12. Statistical analysis

The statistical analysis of the results was performed by using a statistical SPSS/PC program. All values were expressed as mean \pm SE. The results were analyzed using a one-way analysis of variance (ANOVA) test followed by the least significant difference (LSD) test for multiple comparisons. Differences were considered statistically significant at $p < 0.05$.

3. RESULTS

3.1. The effects of fig with olive oil and date-palm fruit extracts (FOD) on the oxidative stress markers of the renal tissue of male rats treated with SiO₂NPs, Al₂O₃NPs, or ZnONPs for 75 days are represented in Table 2

The FOD and FOD-SiO₂NPs-treated groups show insignificant changes in the TAC, GSH, SOD, and TBARS in the renal tissue when compared with the control group.

The SiO₂NPs, Al₂O₃NPs, and ZnONPs-treated groups recorded a significant decrease (P<0.05) in the TAC (18.71%, 31.37%, and 36.17%, respectively), GSH (18.44%, 35.62%, and 41.65%, respectively), SOD (2.59%, 15.12%, and 13.51%, respectively) in contrast to a significant increase (P<0.05) in the TBARS (75.43%, 203.95%, and 208.94%, respectively) in the renal tissue when compared with their corresponding values in the control group.

Similarly, the FOD-Al₂O₃NPs and FOD-ZnONPs-treated groups recorded a significant decrease (P<0.05) in the TAC (19.10%, and 21.61%, respectively), GSH (25.04%, and

31.08%, respectively), SOD (10.00%, and 10.00%, respectively) in contrast to a significant increase (P<0.05) in the TBARS (144.55%, and 152.61%, respectively) in the renal tissue when compared with their corresponding values in the control group.

The FOD-SiO₂NPs, FOD-Al₂O₃NPs, and FOD-ZnONPs-treated groups recorded a significant increase (P<0.05) in the TAC, GSH, and SOD in contrast to a significant decrease (P<0.05) in the TBARS in the renal tissue when compared with their corresponding values in the SiO₂NPs, Al₂O₃NPs, and ZnONPs-treated groups respectively.

Table 2. The effects of fig with olive oil and date-palm fruit extract on the oxidative stress markers of the kidney tissue of male rats treated with silicon oxide nanoparticles, aluminum oxide nanoparticles, or zinc oxide nanoparticles for 75 days

Parameters	Experimental groups							
	Control	FOD	SiO ₂ NPs	FOD-SiO ₂ NPs	Al ₂ O ₃ NPs	FOD-Al ₂ O ₃ NPs	ZnONPs	FOD-ZnONPs
Renal TAC (μmol/g tissue)	2.31±0.04	2.24±0.06	1.88±0.19 ^a	2.16±0.03 ^b	1.58±0.10 ^a	1.87±0.13 ^{a,c}	1.47±0.13 ^a	1.81±0.11 ^{a,d}
		-3.20%	-18.71%	-6.31%	-31.37%	-19.10%	-36.17%	-21.61%
Renal GSH (mmol/g tissue)	5.42±0.06	5.40±0.10	4.42±0.15 ^a	5.24±0.14 ^b	3.49±0.22 ^a	4.06±0.24 ^{a,c}	3.16±0.22 ^a	3.73±0.26 ^{a,d}
		-0.39%	-18.44%	-3.32%	-35.62%	-25.04%	-41.65%	-31.08%
Renal SOD (U/g tissue)	160±0.07	160±0.10	156±1.25 ^a	159±1.28 ^b	136±0.33 ^a	144±1.23 ^{a,c}	138±0.33 ^a	144±1.23 ^{a,d}
		-0.05%	-2.59%	-0.90%	-15.12%	-10.00%	-13.51%	-10.00%
Renal TBARS (nmol/g tissue)	22.6±0.09	22.3±0.43	39.6±2.76 ^a	25.1±1.87 ^b	68.7±1.18 ^a	55.2±2.49 ^{a,c}	69.8±0.37 ^a	57.1±1.92 ^{a,d}
		-1.21%	75.43%	11.04%	203.95%	144.55%	208.94%	152.61%

Note: Results are the mean ± standard error; %, percent of change from the control value; FOD, fig with olive oil and date-palm fruit extracts; SiO₂NPs, silicon oxide nanoparticles; Al₂O₃NPs, aluminum oxide nanoparticles; ZnONPs, zinc oxide nanoparticles; TAC, total antioxidant capacity; GSH, reduced glutathione; SOD, superoxide dismutase activity; TBARS, thiobarbituric acid reactive substances. **For each parameter:** ^ap<0.05, versus control group; ^bp<0.05, versus SiO₂NPs; ^cp<0.05, versus Al₂O₃NPs; ^dp<0.05, versus ZnONPs. **N.B:** The antioxidants (FOD) were used for 2 weeks before and during the administration of the nanoparticles (75 days).

3.2. The effects of fig with olive oil and date-palm fruit extracts (FOD) on the tumor suppressor p53 and inflammatory markers in the renal tissue of male rats treated with SiO₂NPs, Al₂O₃NPs, or ZnONPs for 75 days are represented in Figs. 1-3

The FOD and FOD-SiO₂NPs-treated groups show insignificant changes in the p53, TNF- α , and IL-6 in the renal tissue when compared with control values.

The SiO₂NPs, Al₂O₃NPs, and ZnONPs-treated groups recorded a significant increase ($P < 0.05$) in the p53 (7.87%, 32.96%, and 39.04%, respectively), TNF- α (8.17%, 72.76%, and 132.26%, respectively), and IL-6 (11.77%, 36.65%, and 29.70%, respectively) in the renal tissue when compared with their corresponding values in the control group.




Fig. 1. The effects of fig with olive oil and date-palm fruit extracts (FOD) on the renal tumor necrosis factor- α (TNF- α) in rats treated with different nanoparticles (NPs) for 75 days. ^a $p < 0.05$, versus control group; ^b $p < 0.05$, versus SiO₂NPs; ^c $p < 0.05$, versus Al₂O₃NPs; ^d $p < 0.05$, versus ZnONPs. **N.B:** The antioxidants (FOD) treatments were used for 2 weeks before and during the administration of the nanoparticles.

Similarly, the FOD-Al₂O₃NPs and FOD-ZnONPs-treated groups recorded a significant increase ($P < 0.05$) in the p53 (23.88%, and 25.20%, respectively), TNF- α (50.97%, and 54.66%, respectively), and IL-6 (21.02%, and 17.00%, respectively) in the renal tissue when compared with their corresponding values in the control group.

The FOD-SiO₂NPs, FOD-Al₂O₃NPs, and FOD-ZnONPs-treated groups recorded a significant decrease ($P < 0.05$) in the p53, TNF- α , and IL-6 in the renal tissue when compared with their corresponding values in the SiO₂NPs, Al₂O₃NPs, and ZnONPs-treated groups, respectively.




Fig. 2. The effects of fig with olive oil and date-palm fruit extracts (FOD) on the renal interleukin-6 (IL-6) in rats treated with different nanoparticles (NPs) for 75 days. ^a $p < 0.05$, versus control group; ^b $p < 0.05$, versus SiO₂NPs; ^c $p < 0.05$, versus Al₂O₃NPs; ^d $p < 0.05$, versus ZnONPs. **N.B:** The antioxidants (FOD) treatments were used for 2 weeks before and during the administration of the nanoparticles.




Fig. 3. The effects of fig with olive oil and date-palm fruit extracts (FOD) on the renal tumor suppressor p53 (p53) in rats treated with different nanoparticles (NPs) for 75 days. ^a $p < 0.05$, versus control group; ^b $p < 0.05$, versus SiO₂NPs; ^c $p < 0.05$, versus Al₂O₃NPs; ^d $p < 0.05$, versus ZnONPs. **N.B:** The antioxidants (FOD) treatments were used for 2 weeks before and during the administration of the nanoparticles.

3.3. The effects of fig with olive oil and date-palm fruit extracts (FOD) on the serum kidney function parameters of male rats treated with SiO₂NPs, Al₂O₃NPs, or ZnONPs for 75 days are represented in Table 3

The FOD and FOD-SiO₂NPs-treated groups show insignificant changes in the creatinine, urea, BUN, and uric acid in the serum when compared with control values.

The SiO₂NPs, Al₂O₃NPs, and ZnONPs-treated groups recorded a significant increase (P<0.05) in the creatinine (19.78%, 44.51%, and 62.64%, respectively), urea (23.71%, 43.14%, and 34.57%, respectively), BUN (19.33%, 31.91%, and 43.13%, respectively), and uric acid (10.63%, 19.41%, and 21.80%, respectively) in the renal tissue when compared with their

corresponding values in the control group.

Similarly, the FOD-Al₂O₃NPs and FOD-ZnONPs-treated groups recorded a significant increase (P<0.05) in the creatinine (26.92%, and 44.51%, respectively), urea (24.29%, and 16.00%, respectively), BUN (15.84%, and 25.65%, respectively), and uric acid (8.88%, and 9.95%, respectively) in the renal tissue when compared with their corresponding values in the control group.

The FOD-SiO₂NPs, FOD-Al₂O₃NPs, and FOD-ZnONPs-treated groups recorded a significant decrease (P<0.05) in the creatinine, urea, BUN, and uric acid in the renal tissue when compared with their corresponding values in the SiO₂NPs, Al₂O₃NPs, and ZnONPs-treated groups, respectively.

Table 3. The effects of fig with olive oil and date-palm fruit extract on the serum kidney function parameters of male rats treated with silicon oxide nanoparticles, aluminum oxide nanoparticles, or zinc oxide nanoparticles for 75 days

Parameters	Experimental groups							
	Control	FOD	SiO ₂ NPs	FOD-SiO ₂ NPs	Al ₂ O ₃ NPs	FOD-Al ₂ O ₃ NPs	ZnONPs	FOD-ZnONPs
Serum Creatinine (mg/dL)	0.18±0.00	0.17±0.01	0.21±0.00 ^a	0.19±0.01 ^b	0.26±0.00 ^a	0.23±0.01 ^{a,c}	0.29±0.02 ^a	0.26±0.01 ^{a,d}
		-5.49%	19.78%	8.24%	44.51%	26.92%	62.64%	44.51%
Serum Urea (mg/dL)	35±2.49	34.7±2.27	43.3±1.85 ^a	38.4±2.42 ^b	50.1±2.40 ^a	43.5±1.45 ^{a,c}	47.1±2.23 ^a	40.6±2.30 ^{a,d}
		-0.86%	23.71%	9.71%	43.14%	24.29%	34.57%	16.00%
Serum BUN (mg/dL)	19.1±1.01	19.0±0.94	22.8±1.30 ^a	21.1±1.06 ^b	25.2±1.19 ^a	22.2±0.66 ^{a,c}	27.4±1.16 ^a	24.0±1.37 ^{a,d}
		-0.52%	19.33%	10.20%	31.91%	15.84%	43.13%	25.65%
Serum Uric acid (mg/dL)	2.05±0.08	1.99±0.10	2.26±0.09 ^a	2.14±0.06 ^b	2.44±0.05 ^a	2.23±0.06 ^{a,c}	2.49±0.06 ^a	2.25±0.06 ^{a,d}
		-2.88%	10.63%	4.59%	19.41%	8.88%	21.80%	9.95%

Note: Results are the mean ± standard error; %, percent of change from the control value; FOD, fig with olive oil and date-palm fruit extracts; SiO₂NPs, silicon oxide nanoparticles; Al₂O₃NPs, aluminum oxide nanoparticles; ZnONPs, zinc oxide nanoparticles; BUN, blood urea nitrogen. **For each parameter:** ^ap<0.05, versus control group; ^bp<0.05, versus SiO₂NPs; ^cp<0.05, versus Al₂O₃NPs; ^dp<0.05, versus ZnONPs. **N.B:** The antioxidants (FOD) were used for 2 weeks before and during the administration of the nanoparticles (75 days).

3.4. The effects of fig with olive oil and date-palm fruit extracts (FOD) on the kidney histopathological characters of male rats treated with SiO₂NPs, Al₂O₃NPs, or ZnONPs for 75 days

The control kidney showed a normal morphological appearance. However, the antioxidant-treated NPs-administered groups

(FOD-SiO₂NPs, FOD-Al₂O₃NPs, and FOD-ZnONPs) recorded significantly ameliorated histopathological characters in the kidney tissue when compared with their corresponding values in the non-antioxidant-treated NPs-administered groups (SiO₂NPs, Al₂O₃NPs, and ZnONPs, respectively) as indicated in (Table 4 and Fig. 4).

Table 4. The effects of fig with olive oil and date-palm fruit extract on the kidney histopathological results of male rats treated with silicon oxide nanoparticles, aluminum oxide nanoparticles, or zinc oxide nanoparticles for 75 days

	G	BS	Tubules			Interstitium	Medulla
			Lining	Brush border	Lumen		
Control	0	0	0	0	0	0	0
SiO ₂ NPs	+	+	+	0	+	+	+
FOD-SiO ₂ NPs	0	0	0	0	0	+	0
Al ₂ O ₃ NPs	+	+	+	0	0	+	+
FOD-Al ₂ O ₃ NPs	0	0	+	0	0	+	+
ZnONPs	++	+	++	++	++	++	+
FOD-ZnONPs	0	0	+	0	+	+	+

FOD, fig with olive oil and date-palm fruit extracts; SiO₂NPs, silicon oxide nanoparticles; Al₂O₃NPs, aluminum oxide nanoparticles; ZnONPs, zinc oxide nanoparticles.

Glomeruli (G): (0, average; +, edematous/congested; ++, small-sized/atrophied).

Bowman's spaces (BS): (0, average; +, widened/dilated; ++, obliterated).

Tubules

- Lining: (0, average; +, edematous/apoptotic; ++, atrophied/necrotic).
- Brush border: (0, preserved; +, partial loss; ++, complete loss).
- Lumen: (0, free; +, few/scattered casts; ++, marked intra-tubular casts).

Interstitium: (0, average; +, dilated/congested BV; ++, markedly dilated BV/interstitial hemorrhage).

Medulla: (0, average; +, congested capillaries/scattered hyaline casts; ++, marked hyaline casts).




Fig. 4. The effects of fig with olive oil and date-palm fruit extracts (FOD) on the microscopic histological appearance of livers treated with different nanoparticles (NPs) for 75 days. **A**, higher power view of the control kidney showing average glomeruli (G) with average Bowman's spaces (BS), average proximal tubules (P) with preserved brush borders (black arrows), average distal tubules (D), and average interstitium (blue arrow); **B**, FOD-treated group renal medulla showing average collecting tubules (CT) with average epithelial lining (black arrow) and average peri-tubular capillaries (blue arrow); **C**, SiO₂NPs-treated group showing atrophied glomerulus (G) with widened Bowman's space (BS), proximal tubules (P) with mildly edematous epithelial lining (black arrow), intra-tubular hyaline casts (blue arrow), and mildly congested blood vessels (red arrow); **D**, FOD-SiO₂NPs-treated group showing average glomeruli (G) with average Bowman's spaces (BS), proximal tubules (P) with average epithelial lining (black arrow), and average interstitium (blue arrow); **E**, Al₂O₃ NPs-treated group showing average glomerulus (G) with average Bowman's space (BS), proximal tubules (P) with mildly edematous epithelial lining (black arrow) and mildly congested interstitial blood vessels (blue arrow); **F**, FOD-Al₂O₃NPs-treated group showing average glomerulus (G) with average Bowman's space (BS), proximal tubules (P) with mildly edematous (black arrow) and apoptotic epithelial lining (blue arrow); **G**, ZnO NPs-treated group showing atrophied glomerulus (G) with widened Bowman's space (BS), proximal tubules (P) with edematous epithelial lining (black arrow), and areas of interstitial hemorrhage (Hge) with mild inflammatory infiltrate (blue arrow); **H**, FOD-ZnO NPs-treated group showing average glomerulus (G) with average Bowman's space (BS), proximal tubules (P) with mildly edematous (black arrow) and apoptotic epithelial lining (blue arrow).

4. DISCUSSION

Recently, more nanomedicines have been already approved by the U.S. Food and Drug Administration (FDA) for human use [29]. With the progress in the nanotechnology field, there may be an increase in the exposure of humans to nanoparticles, so further urgent studies are required to study the possibility of any detrimental health impacts, target organs damage as well as their mechanisms [30].

The present study confirmed the renal toxicity of rats treated with SiO₂NPs, Al₂O₃NPs, or ZnONPs [31-34] which indicated that these toxicities may be mediated through induction of oxidative stress, and lipid peroxidation, systemic inflammation, and renal toxicity. Due to their small size, the NPs can travel to different organs of the body when ingested. The mechanism(s) of the kidney cytotoxicity induced by nanoparticles might be due to enhanced production of reactive oxygen species ROS leading to oxidative stress and lipid peroxidation and induction of inflammatory pathways [3, 35].

Regarding the oxidative stress markers, the SiO₂NPs, Al₂O₃NPs, and ZnONPs-treated groups recorded a significant depletion of TAC, GSH, SOD, and CAT and a significant increase of TBARS as compared to the control group. These results are in agreement with previous studies [22, 32, 34, 36] which reported a significant depletion in the antioxidant system accompanied by a dramatic fall in the antioxidants after SiO₂NPs, Al₂O₃NPs, or ZnONPs administration.

The lipid peroxidation products (TBARS) are correlated with increased production of ROS so, considered a useful marker of oxidative stress. In addition, TBARS played an important role in determining the NPs (SiO₂NPs, Al₂O₃NPs, or ZnONPs) toxicity [32, 36]. The cells are equipped with several natural enzymatic (SOD and CAT) and non-enzymatic (GSH) endogenous

antioxidant defenses to combat oxidative stress [37]. The treatments with SiO₂NPs, Al₂O₃NPs, or ZnONPs lead to the depletion of these endogenous antioxidants [22, 32]. In agreement, Guan *et al.* [38] and Zhang *et al.* [3] recorded that SiO₂NPs, Al₂O₃NPs, or ZnONPs administration cause oxidative stress leading to DNA damage and cytotoxicity in the kidney.

The results obtained in the present study showed that the extra-virgin olive oil with fig and date-palm fruit extracts are quite safe products and they have not resulted in any adverse effects when administered to rats [39-43].

The present study demonstrated that the antioxidant-treated NPs-administered groups (FOD-SiO₂NPs, FOD-Al₂O₃NPs, and FOD-ZnONPs) recorded significantly ameliorated oxidative stress markers in the kidney tissue when compared with their corresponding values in the non-antioxidant-treated NPs-administered groups (SiO₂NPs, Al₂O₃NPs, and ZnONPs, respectively). The treatment with FOD revealed the presence of numerous and varied antioxidants and those diverse natural antioxidants acted synergistically [40, 44]. The free radical toxicity in the present study was counteracted by the FOD treatment which might be attributed to the protective activity of the plant antioxidants since they are complex mixtures of many chemicals [45].

The SiO₂NPs, Al₂O₃NPs, and ZnONPs-treated groups recorded a significant increase in the renal TNF- α and renal IL-6 when compared with their corresponding control values which revealed the inflammatory reaction appeared after rats exposed to these NPs as agreeing with [46] who concluded that the pro-inflammatory cytokines (TNF- α , IL-6, and IL-1 β) could play an important role in regulating immunity. In agreement, Faddah *et al.* [47] found that the elevation in serum inflammatory markers including TNF- α , IL-6, and CRP due to NPs

ingestion induced nephrotoxicity.

The SiO₂NPs, Al₂O₃NPs, and ZnONPs-treated groups recorded a significant increase in the p53 when compared with their corresponding control values. It is agreeing with the fact that oxidative stress can stimulate apoptotic cell death [48]. Tumor suppressor genes regulate diverse cellular activities including programmed cell death. In addition, p53 known as a potent prooxidant protein, down-regulates intracellular ROS levels, thus reducing the probability of genetic alterations and apoptosis [49, 50]. In agreement, different NPs such as SiO₂NPs or ZnONPs caused DNA damage and apoptosis in the kidney tissues [32, 51].

Our results showed that FOD effectively reduced inflammatory markers and p53 induced by the NPs-administration. These results revealed that FOD contains a strong antioxidant effect, and since they contained high amounts of phenolic compounds, they have been reported as free radical scavengers and have therapeutic effects against tissue inflammatory, cancer (tumor suppressor), and cardiovascular diseases [40, 52, 53].

The kidney function parameters of urea, creatinine and uric acid levels were markedly increased by SiO₂NPs, Al₂O₃NPs, or ZnONPs administration in the present study. In kidney damage, these biomarkers were released into the bloodstream from the proximal cells. In agreement, the elevated concentration of these biomarkers indicates proximal cell destruction [54], and the inflammatory cells were infiltrated into the renal tissue after oral ingestion of SiO₂NPs, Al₂O₃NPs, or ZnONPs in rats [55]. Also, in agreement, Yan *et al.* [51] and Yousef *et al.* [22] demonstrated that ZnONPs and Al₂O₃NPs exhibited toxicological symptoms implying the potential renal damage. In addition, the *in vitro* research indicated the NP's cytotoxicity to kidney cells [56, 57].

The most sensitive biochemical markers in the diagnosis of renal damage are the retention of creatinine and urea in the blood [58]; therefore, the significant increase in these biomarkers indicates kidney functional damage [59]. The higher improvement in the antioxidant markers by concomitant administration of FOD in the present study could be explained by Allouche *et al.* [60] who stated that olive oil by its richness in carotenoids, tocopherols, vitamin E, and polyphenols as potential antioxidants properties acting as ROS scavenger as well as increasing the activity of antioxidant enzymes. In agreement, the hydroxytyrosol of FOD was reported to be an effective scavenger. Administration of hydroxytyrosol to the albino rats prevented kidney toxicity [61]. The decrease in uric acid level may be due to FOD treatment that enhances GSH concentration and inhibits XDH conversion to XO that catalyzes purine degradation, hypoxanthine, and xanthine metabolism to uric acid [40].

The kidney damage caused by SiO₂NPs, Al₂O₃NPs, or ZnONPs in the present study is further confirmed by histopathological examination, which showed many pathological features such as degeneration of the epithelial cells of various renal tubules, distorted corpuscles, capillaries shrinkage, and inflammatory infiltration in various tubules. These data are in line with the previous studies [22, 62, 63].

The significantly ameliorated histopathological characters recorded at the FOD-SiO₂NPs, FOD-Al₂O₃NPs, and FOD-ZnONPs when compared with their corresponding values in the non-antioxidant-treated NPs groups could be attributed to the antioxidant effect of FOD due to their varied phenolic compounds, which were able to donate a hydrogen atom to the free radicals, thus stopping the propagation chain reaction during the lipid peroxidation process

[40, 64].

Conclusion

According to the results obtained in the present study, the administration of extra virgin olive oil with fig and date palm extracts (FOD) provides considerable protective effects against different NPs-induced subchronic nephrotoxicity in male Wistar albino rats. In addition, the administration of FOD revealed a synergistic effect of the combination between them to produce a broad spectrum of antioxidative activities that creates an effective defense system against the free radical attack.

Declarations

Ethics approval and consent to participate

The study protocol was approved by the Research Ethical Committee, Faculty of Pharmacy, Ain Shams University, and conducted according to the regulations and recommendations of the ethical guidelines and complied with the guide for the care and use of laboratory animals.

Consent to publish

All authors have read and agreed to the published version of the manuscript.

Availability of data and materials

Data analyzed during this study are all included in the main manuscript.

Competing interests

No competing interests were declared by the authors.

Funding statement

No funding source was received

5. REFERENCES

1. Oberdörster, G., Oberdörster, E., and Oberdörster, J., Nanotoxicology: an emerging discipline evolving from studies of ultrafine

particles. Environmental health perspectives, 2005. 113(7): p. 823-839.

2. Dağlıoğlu, Y. and Yılmaz Öztürk, B., Effect of concentration and exposure time of ZnO-TiO₂ nanocomposite on photosynthetic pigment contents, ROS production ability, and bioaccumulation of freshwater algae (*Desmodesmus multivariable*). *Caryologia*, 2018. 71(1): p. 13-23.
3. Zhang, Y., Bai, Y., Jia, J., Gao, N., Li, Y., Zhang, R., Jiang, G., and Yan, B., Perturbation of physiological systems by nanoparticles. *Chemical Society Reviews*, 2014. 43(10): p. 3762-3809.
4. Xu, Y., Wang, N., Yu, Y., Li, Y., Li, Y.-B., Yu, Y.-B., Zhou, X.-Q., and Sun, Z.-W., Exposure to silica nanoparticles causes reversible damage of the spermatogenic process in mice. *PloS one*, 2014. 9(7): p. e101572.
5. Nemmar, A., Yuvaraju, P., Beegam, S., Yasin, J., Kazzam, E. E., and Ali, B. H., Oxidative stress, inflammation, and DNA damage in multiple organs of mice acutely exposed to amorphous silica nanoparticles. *International journal of nanomedicine*, 2016. 11: p. 919.
6. El-Adawy, M., Abd El-Aziz, M., El-Shazly, K., Ali, N. G., and El-Magd, M. A., Dietary propionic acid enhances antibacterial and immunomodulatory effects of oxytetracycline on Nile tilapia, *Oreochromis niloticus*. *Environmental Science and Pollution Research*, 2018. 25(34): p. 34200-34211.
7. Abdelwahab, M., El-Barbary, A., El-Said, K., Betiha, M., Elkholy, H., Chiellini, E., and El-Magd, M., Functionalization of poly (3-hydroxybutyrate) with different thiol compounds inhibits MDM2-p53 interactions in MCF7 cells. *Journal of Applied Polymer Science*, 2019. 136(2): p. 46924.

8. Wagner, A. J., Bleckmann, C. A., Murdock, R. C., Schrand, A. M., Schlager, J. J., and Hussain, S. M., Cellular interaction of different forms of aluminum nanoparticles in rat alveolar macrophages. *The journal of physical chemistry B*, 2007. 111(25): p. 7353-7359.
9. Rastogi, A., Zivcak, M., Sytar, O., Kalaji, H. M., He, X., Mbarki, S., and Brestic, M., Impact of metal and metal oxide nanoparticles on the plant: a critical review. *Frontiers in Chemistry*, 2017. 5: p. 78.
10. Li, M., Lin, D., and Zhu, L., Effects of water chemistry on the dissolution of ZnO nanoparticles and their toxicity to *Escherichia coli*. *Environmental pollution*, 2013. 173: p. 97-102.
11. Espitia, P. J. P., Soares, N. d. F. F., dos Reis Coimbra, J. S., de Andrade, N. J., Cruz, R. S., and Medeiros, E. A. A., Zinc oxide nanoparticles: synthesis, antimicrobial activity, and food packaging applications. *Food and bioprocess technology*, 2012. 5(5): p. 1447-1464.
12. Hackenberg, S., Zimmermann, F. Z., Scherzed, A., Friehs, G., Froelich, K., Ginzkey, C., Koehler, C., Burghartz, M., Hagen, R., and Kleinsasser, N., Repetitive exposure to zinc oxide nanoparticles induces DNA damage in human nasal mucosa mini organ cultures. *Environmental and molecular mutagenesis*, 2011. 52(7): p. 582-589.
13. Tsigalou, C., Konstantinidis, T., Paraschaki, A., Stavropoulou, E., Voidarou, C., and Bezirtzoglou, E., Mediterranean diet as a tool to combat inflammation and chronic diseases. An overview. *Biomedicine*, 2020. 8(7): p. 201.
14. Solomon, A., Golubowicz, S., Yablowicz, Z., Grossman, S., Bergman, M., Gottlieb, H. E., Altman, A., Kerem, Z., and Flaishman, M. A., Antioxidant activities and anthocyanin content of fresh fruits of common fig (*Ficus carica L.*). *J. Agric. Food Chem.*, 2006. 54(20): p. 7717-7723.
15. Lansky, E. P., Paavilainen, H. M., Pawlus, A. D., and Newman, R. A., *Ficus spp.* (fig): Ethnobotany and potential as anticancer and anti-inflammatory agents. *J. Ethnopharmacol.*, 2008. 119: p. 195-213.
16. Quiles, J. L., Ramírez-Tortosa, M. C., and Yaqoob, P., *Olive oil and health*, Cambridge MA: Cabi International. 2006.
17. Parvin, S., Easmin, D., Sheikh, A., Biswas, M., Sharma, S. C. D., Jahan, M. G. S., Islam, M. A., Shovon, M., and Roy, N., Nutritional analysis of date fruits (*Phoenix dactylifera L.*) in perspective of Bangladesh. *American Journal of Life Sciences*, 2015. 3(4): p. 274-278.
18. Al-Alawi, R. A., Al-Mashiqri, J. H., Al-Nadabi, J. S., Al-Shihi, B. I., and Baqi, Y., Date palm tree (*Phoenix dactylifera L.*): natural products and therapeutic options. *Frontiers in plant science*, 2017. 8: p. 845.
19. Siddiq, M. and Greiby, I., Overview of date fruit production, postharvest handling, processing, and nutrition. *Dates: postharvest science, processing technology, and health benefits*. 1st edition. Edited by M. Siddiq, SM Aleid, and AA Kader. John Wiley and Sons, Chichester, United Kingdom, 2014: p. 1-28.
20. Gmoshinski, I. V., Shipelin, V. A., Shumakova, A. A., Trushina, E. N., Mustafina, O. K., Safenkova, I. V., Khotimchenko, S. A., Nikityuk, D. B., and Tutelyan, V. A., Toxicity Evaluation of Nanostructured Silica Orally Administered to Rats: Influence on Immune System Function. *Nanomaterials*, 2020. 10(11): p. 2126.

21. Pakrashi, S., Dalai, S., Sabat, D., Singh, S., Chandrasekaran, N., and Mukherjee, A., Cytotoxicity of Al₂O₃ nanoparticles at low exposure levels to a freshwater bacterial isolate. *Chemical research in toxicology*, 2011. 24(11): p. 1899-1904.
22. Yousef, M. I., Mutar, T. F., and Kamel, M. A. E.-N., Hepato-renal toxicity of oral sub-chronic exposure to aluminum oxide and/or zinc oxide nanoparticles in rats. *Toxicology reports*, 2019. 6: p. 336-346.
23. Gilani, A. H., Mehmood, M. H., Janbaz, K. H., Khan, A. U., and Saeed, S. A., Ethnopharmacological studies on antispasmodic and antiplatelet activities of *Ficus carica*. *J. Ethnopharmacol.*, 2008. 119(1): p. 1-5.
24. Al-Qarawi, A. A., Abdel-Rahman, H., Ali, B. H., Mousa, H. M., and El-Mougy, S. A., The ameliorative effect of dates (*Phoenix dactylifera* L.) on ethanol-induced gastric ulcer in rats. *J. Ethnopharmacol.*, 2005. 98(3): p. 313-317.
25. Fathy, A. H., Bashandy, M. A., Bashandy, S. A., Mansour, A. M., and Azab, K. S., The beneficial effect of natural antioxidants from olive oil with fig and date palm fruit extracts on biochemical and hematological parameters in rats treated with doxorubicin and γ -radiation. *FACETS*, 2018. 3(1): p. 722-735.
26. Vinson, J. A., Zubik, L., Bose, P., Samman, N., and Proch, J., Dried fruits: excellent in vitro and in vivo antioxidants. *J. Am. Coll. Nutr.*, 2005. 24(1): p. 44-50.
27. Institute of Laboratory Animal Resources, Guide for the Care and Use of Laboratory Animals, eighth edition. Committee for the update of the guide and use of laboratory animals. National research council of the national academies. National Academy Press. 1996: Washington, D.C.
28. Bancroft, J. D., and Gamble, M., editors. *Theory and practice of histological techniques*. Elsevier health sciences, 2008.
29. Bobo, D., Robinson, K. J., Islam, J., Thurecht, K. J., and Corrie, S. R., Nanoparticle-based medicines: a review of FDA-approved materials and clinical trials to date. *Pharmaceutical research*, 2016. 33(10): p. 2373-2387.
30. Yao, Y., Zang, Y., Qu, J., Tang, M., and Zhang, T., The toxicity of metallic nanoparticles on the liver: the subcellular damages, mechanisms, and outcomes. *International journal of nanomedicine*, 2019. 14: p. 8787.
31. Sharma, V., Singh, P., Pandey, A. K., and Dhawan, A., Induction of oxidative stress, DNA damage and apoptosis in mouse liver after sub-acute oral exposure to zinc oxide nanoparticles. *Mutation Research/Genetic Toxicology and Environmental Mutagenesis*, 2012. 745(1-2): p. 84-91.
32. Du, Z., Zhao, D., Jing, L., Cui, G., Jin, M., Li, Y., Liu, X., Liu, Y., Du, H., and Guo, C., Cardiovascular toxicity of different sizes amorphous silica nanoparticles in rats after intratracheal instillation. *Cardiovascular toxicology*, 2013. 13(3): p. 194-207.
33. He, X., In vivo nanotoxicity assays in animal models. *Toxicology of Nanomaterials*, 2016: p. 151-198.
34. Demir, E., A review on nanotoxicity and nanogenotoxicity of different shapes of nanomaterials. *Journal of Applied Toxicology*, 2021. 41(1): p. 118-147.
35. Hillyer, J. F. and Albrecht, R. M., Gastrointestinal persorption and tissue distribution of differently sized colloidal gold nanoparticles. *Journal of pharmaceutical sciences*, 2001. 90(12): p. 1927-1936.

36. Wu, T. and Tang, M., Review of the effects of manufactured nanoparticles on mammalian target organs. *Journal of Applied Toxicology*, 2018. 38(1): p. 25-40.
37. Halliwell, B., Gutteridge, J. M., and Cross, C. E., Free radicals, antioxidants, and human disease: where are we now? *J. Lab. Clin. Med.*, 1992. 119(6): p. 598-620.
38. Guan, R., Kang, T., Lu, F., Zhang, Z., Shen, H., and Liu, M., Cytotoxicity, oxidative stress, and genotoxicity in human hepatocyte and embryonic kidney cells exposed to ZnO nanoparticles. *Nanoscale research letters*, 2012. 7(1): p. 1-7.
39. Bashandy, M. A., Abd-el-aal, A., Ibrahim, D. F., and El-sharkawy, M. A., Protective effects of date palm extract as natural antioxidants on hepatotoxicity induced by *Cerastes cerastes* venom in albino rats. *Int. J. Adv. Res.*, 2016. 4: p. 647-665.
40. Bashandy, M. A., Abd El- Rasheid, H. G., Hasan, H. F., and Fathy, A. H., Protective and therapeutic effects of olive oil and *Ficus carica* as natural antioxidants on some biochemical parameters in liver of γ -irradiated male albino rats *Al Azhar Bulletin of science*, 2014. 25(1): p. 1-16.
41. Sruthi, B., Sunny, G., Hajra, N., and Sakthivel, S., Diuretic activity of ethanolic extracts of fruits. *Int. J. Res. Pharmacol. Pharmacotherapy.*, 2012. 1(1): p. 25-28.
42. Sheikh, B. Y., Elsayed, W. M., Samman, A. H., Sheikh, B. Y., and Ladin, A. M., Ajwa dates as a protective agent against liver toxicity in rats. *Eur. Sci. J.*, 2014. 3: p. 358-368.
43. Zhang, J., Zhu, W.-f., Xu, J., Kitdamrongtham, W., Manosroi, A., Manosroi, J., Tokuda, H., Abe, M., Akihisa, T., and Feng, F., Potential cancer chemopreventive and anticancer constituents from the fruits of *Ficus hispida* Lf (Moraceae). *J. Ethnopharmacol.*, 2018. 214: p. 37-46.
44. Viola, P. and Viola, M., Virgin olive oil as a fundamental nutritional component and skin protector. *Clin. Dermatol.*, 2009. 27: p. 159-165.
45. Jagetia, G. C., Radioprotective potential of plants and herbs against the effects of ionizing radiation. *J. Clin. Biochem. Nutr.*, 2007. 40: p. 74-81.
46. Libby, P., Ridker, P. M., and Maseri, A., Inflammation, and atherosclerosis. *Circulation*, 2002. 105(9): p. 1135-1143.
47. Faddah, L. M., Baky, N. A. A., Al-Rasheed, N. M., Al-Rasheed, N. M., Fatani, A. J., and Atteya, M., Role of quercetin and arginine in ameliorating nano zinc oxide-induced nephrotoxicity in rats. *BMC complementary and alternative medicine*, 2012. 12(1): p. 1-14.
48. Chang, M.-C., Chan, C.-P., Wang, Y.-J., Lee, P.-H., Chen, L.-I., Tsai, Y.-L., Lin, B.-R., Wang, Y.-L., and Jeng, J.-H., Induction of necrosis and apoptosis to KB cancer cells by sanguinarine is associated with reactive oxygen species production and mitochondrial membrane depolarization. *Toxicology and applied pharmacology*, 2007. 218(2): p. 143-151.
49. Achanta, G. and Huang, P., Role of p53 in sensing oxidative DNA damage in response to reactive oxygen species-generating agents. *Cancer Research*, 2004. 64(17): p. 6233-6239.
50. Vurusaner, B., Poli, G., and Basaga, H., Tumor suppressor genes and ROS: complex networks of interactions. *Free Radical Biology and Medicine*, 2012. 52(1): p. 7-18.
51. Yan, G., Huang, Y., Bu, Q., Lv, L., Deng, P., Zhou, J., Wang, Y., Yang, Y., Liu, Q., and Cen, X., Zinc oxide nanoparticles cause

- nephrotoxicity and kidney metabolism alterations in rats. *Journal of Environmental Science and Health, Part A*, 2012. 47(4): p. 577-588.
52. Cai, L. and Cherian, M. G., Zinc-metallothionein protect from DNA damage induced by radiation better than glutathione and copper or cadmium-metallothionein. *Toxicol. Lett.*, 2003. 136(3): p. 193-198.
53. Magrone, T., Effects of extra virgin olive oil on the immune-mediated inflammatory responses: Potential clinical applications. *Endocr. Metab. Immune. Disord. Drug Targets*, 2018. 18(1): p. 3-3.
54. Zhu, H.-c. and Cao, R.-l., The relationship between serum levels of uric acid and prognosis of infection in critically ill patients. *World journal of emergency medicine*, 2012. 3(3): p. 186.
55. Chien, C.-C., Yan, Y.-H., Juan, H.-T., Cheng, T.-J., Liao, J.-B., Lee, H.-P., and Wang, J.-S., Sustained renal inflammation following 2 weeks of inhalation of occupationally relevant levels of zinc oxide nanoparticles in Sprague Dawley rats. *Journal of toxicologic pathology*, 2017. 30(4): p. 307-314.
56. Kim, J. S., Kuk, E., Yu, K. N., Kim, J.-H., Park, S. J., Lee, H. J., Kim, S. H., Park, Y. K., Park, Y. H., and Hwang, C.-Y., Antimicrobial effects of silver nanoparticles. *Nanomedicine: Nanotechnology, biology and medicine*, 2007. 3(1): p. 95-101.
57. Nagesha, D. K., Whitehead, M. A., and Coffer, J. L., Biorelevant calcification and non-cytotoxic behavior in silicon nanowires. *Advanced Materials*, 2005. 17(7): p. 921-924.
58. Nwanjo, H. U., Efficacy of aqueous leaf extract of *Vernonia amygdalina* on plasma lipoprotein and oxidative status in diabetic rat models. *Nigerian Journal of Physiological Sciences*, 2005. 20(1): p. 39-42.
59. Panda, P. R., Dutt, N. D., and Nicolau, A., Memory issues in embedded systems-on-chip: optimizations and exploration. 1999: Springer Science & Business Media.
60. Allouche, Y., Beltrán, G., Gaforio, J. J., Uceda, M., and Mesa, M. D., Antioxidant and antiatherogenic activities of pentacyclic triterpenic diols and acids. *Food and Chemical Toxicology*, 2010. 48(10): p. 2885-2890.
61. Eliza, J., Daisy, P., Ignacimuthu, S., and Duraipandiyan, V., Antidiabetic and antilipidemic effect of eremanthin from *Costus speciosus* (Koen.) Sm., in STZ-induced diabetic rats. *Chemico-Biological Interactions*, 2009. 182(1): p. 67-72.
62. Xia, T., Kovochich, M., Liong, M., Madler, L., Gilbert, B., Shi, H., Yeh, J. I., Zink, J. I., and Nel, A. E., Comparison of the mechanism of toxicity of zinc oxide and cerium oxide nanoparticles based on dissolution and oxidative stress properties. *ACS nano*, 2008. 2(10): p. 2121-2134.
63. Saman, S., Moradhaseli, S., Shokouhian, A., and Ghorbani, M., Histopathological effects of ZnO nanoparticles on liver and heart tissues in Wistar rats. *Adv Biores*, 2013. 4(2): p. 83-8.
64. Yanishlieva, N. V. and Marinova, E. M., Activity and mechanism of action and a natural antioxidant in lipids. *Recent. Res. Develop.*, 1998. 2: p. 1-14.